

Single Nuclei Preparation for Mouse Kidney

This protocol was used in

[5 Million Mouse Single-Cell Atlas from 7 Tissues](#)

Materials

Material	Supplier	Part Number
Sucrose	Sigma-Aldrich [®]	S0389
KCl	Invitrogen [®]	AM9640G
MgCl ₂	Invitrogen	AM9530g
Tris buffer, pH 8	Invitrogen	AM9856
Nuclease-Free Water	Sigma-Aldrich	W4502
DTT	Invitrogen	P2325
RNase Inhibitor	QIAGEN [®]	Y9240
SUPERase-In RNase Inhibitor	Invitrogen	AM2696
Triton X-100	Sigma-Aldrich	T8787
DNA low binding, DNase/RNase free, filtered pipette tips 20 µL, 200 µL, 1,000 µL	Varies	Varies
Falcon [®] High Clarity PP Centrifuge Tubes, 15 mL	Corning [®]	352097
5.0mL LoBind [®] Tubes	Eppendorf [®]	0030108302

Corning 100 mm TC-treated Culture Dish	Corning	430167
RNase-free razor blade	Varies	Varies
Cell Strainer (40 µm)	Corning	431750
Trypan Blue	Varies	Varies
RNaseZAP [®] RNase Decontamination Solution	Thermo Fisher Scientific [®]	AM9780
Ethyl alcohol, Pure	Sigma-Aldrich	459844
Optional: Fluorescent stain (e.g., Hoechst, DAPI, or AOPI)	Varies	Varies
Dounce Tissue Grinder, 1mL	WHEATON [®]	357538
Single channel pipettes: P20, P200, P1000	Varies	Varies
Cold block	Varies	Varies
RNase-free forceps	Varies	Varies
Hemocytometer	Sigma-Aldrich	Z359629



Note: In addition to the above materials, you will need access to dry ice for this protocol.

Preparation

1. Clean the bench top and dissection area with 70% ethanol, followed by RNaseZAP to remove RNases.

2. Prepare the following solutions, and add reagents in the order provided in the table.
3. Clean the Dounce homogenizer with RNaseZAP followed by 70% ethanol then rinse well with dH₂O and let dry.

NIM1 buffer

Reagents	Stock Concentration	Final Concentration	Volume for 15 mL
Sucrose	1.5 M	250 mM	2,500 µL
KCl	2 M	25 mM	187.5 µL
MgCl ₂	1 M	5 mM	75 µL
Tris buffer, pH 8	1 M	10 mM	150 µL
Water	-	-	12,087.5 µL

Homogenization buffer

Reagents	Stock Concentration	Final Concentration	Volume for 5 mL
NIM1 Buffer	1.5 M		4,845 µL
DTT	1 mM	1 µM	5 µL
RNase Inhibitor	40 U/µL	0.4U/µL	50 µL
SUPERase-In RNase Inhibitor	20 U/µL	0.2U/µL	50 µL
Triton X-100	10%	-	50 µL



Note: If there is a potential that your sample was exposed to any external sources of RNases during tissue harvest or if your tissue is known to have elevated RNase activity, you can optionally add in additional RNase inhibitors of your choosing when preparing the homogenization buffer.

Nuclei Extraction

1. Gather a dounce and keep at 4°C until use.
2. Make the NIM1 buffer and keep on ice until use.
3. Make the homogenization buffer and keep on ice until use.
4. Prior to douncing, fill a bucket with dry ice and place a cold block on top of the dry ice to use as a cold surface. Place a culture plate on top of the cold block and wait 5 minutes so the culture plate is sufficiently cooled.
5. Transfer the kidney from the cryovial onto the pre-cooled culture plate. Using RNase-free forceps and a razor blade, make a longitudinal cut to separate the kidney into two halves. Slice each half into 0.5 cm slices.
6. Dounce
 - a. Add tissue sample to the dounce and resuspend in 700 μ L of homogenization buffer.
 - b. Perform 5 strokes of loose pestle.
 - c. Perform up to 10 - 15 strokes of tight pestle. Add additional strokes of the tight pestle if you still see large pieces.
 - d. Add additional homogenization buffer up to 1.4 mL.

7. Filter homogenates with a 40 μ m strainer into a 5mL or 15 mL tube.
8. Count the nuclei using a preferred accurate method.



Note: In our experience, nuclei are much stickier than whole cells.

For researchers making the single-nuclei suspension for the first time, we suggest confirming that the sample contains mostly single nuclei. To do this, take 10 μ L of the sample and stain it with Hoechst, DAPI, or AOPI for more than 5 min, place it on a hemocytometer, and visualize it under a 20x and 40x objective on an epifluorescence microscope. A best practices resource for hemocytometer use, see: <https://www.hemocytometer.org/>

9. Move up to the maximum number of nuclei* allowed by the Evercode™ Fixation kit into the nuclei fixation workflow.

*Up to 1 million nuclei per reaction for [Evercode Nuclei Fixation v3](#) and up to 100,000 nuclei for [Evercode Low Input Nuclei Fixation v3](#).

Notice

This document and its contents may be proprietary to Parse Biosciences, Inc. ("Parse Biosciences") and is intended solely for use by its customers in connection with the use of the product(s) described herein and for no other purpose. The products may be used solely FOR RESEARCH PURPOSES, AND MAY NOT BE USED IN ANY DIAGNOSTIC OR THERAPEUTIC USE IN HUMANS OR ANIMALS. This document and its contents shall not be used or distributed for any other purpose and/ or otherwise communicated, disclosed or reproduced in any way whatsoever without the prior written consent of Parse Biosciences. No rights are granted under this document with respect to any of Parse Biosciences' intellectual property rights. The license to use of any products described herein is subject to a separate written agreement between Parse Biosciences and the applicable user. Parse Biosciences shall have no liability for any direct, indirect, consequential or incidental damages arising out of any failure to use the product(s) in strict compliance with the terms herein. This document may contain references to third-party sources of information, hardware or software, products, or services, or intellectual property including but not limited to trademarks, and/ or third-party web sites (collectively "Third Party Information and Intellectual Property"). Parse Biosciences does not control, and is not responsible for, any Third Party Information or Intellectual Property. The inclusion of Third Party Information or Intellectual Property in this document does not imply endorsement by Parse Biosciences of the Third Party Information or Intellectual Property or the third party in any way. A non-exhaustive list of Parse Biosciences trademarks can be found at www.parsebiosciences.com/legal/trademarks/. Parse Biosciences products may be covered by one or more patents as listed at www.parsebiosciences.com/legal/patents/. The product(s) offer by Parse Biosciences described in this document are provided for one-time use by the purchaser and may not be re-used, refurbished or resold. In addition, such product(s) may not be altered, changed or modified by anyone other than Parse Biosciences and its authorized agents, and Parse Biosciences will not be liable for any such alterations, changes or modifications.